

Appendix C | **Listed Species in the County of Los Angeles**

Appendix C | Listed Species in the County of Los Angeles

LISTED PLANT SPECIES WITH POTENTIAL TO OCCUR IN THE COUNTY OF LOS ANGELES (CDFG 2010)

Scientific Name Common Name	Special Status	Preferred Habitat
<i>Acmispon argophyllus</i> var. <i>adsurgens</i> San Clemente Island bird's-foot trefoil	SE, 1B	Rocky volcanic substrates with coastal scrub and coastal bluff scrub (15-395 meters)
<i>Acmispon dendroideus</i> var. <i>traskiae</i> San Clemente Island lotus	FE, SE, 1B	Coastal scrub, coastal bluff scrub, valley and foothill grassland (15-365 meters)
<i>Arenaria paludicola</i> Marsh sandwort	FE, SE, 1B	Marshes and swamps (10-170 meters)
<i>Astragalus brauntonii</i> Braunton's milk-vetch	FE, 1B	Gravelly clay soils in closed-cone coniferous forest, chaparral, coastal scrub, valley and foothill grasslands (4-640 meters)
<i>Astragalus pycnostachyus</i> var. <i>lanosissimus</i> Ventura Marsh milk-vetch	FE, SE, 1B	Coastal salt marsh (1-35 meters)
<i>Astragalus tener</i> var. <i>titi</i> Coastal dunes milk-vetch	FE, SE, 1B	Moist, sandy depressions in coastal bluff scrub, coastal dunes (1-50 meters)
<i>Berberis nevini</i> Nevin's barberry	FE, SE, 1B	Chaparral, cismontane woodland, coastal scrub, riparian scrub (290-1,575 meters)
<i>Brodiaea filifolia</i> Thread-leaved brodiaea	FT, SE, 1B	Cismontane woodland, coastal scrub, playas, valley and foothill grassland, vernal pools (25-860 meters)
<i>Castilleja gleasoni</i> Mt. Gleason paintbrush	1B	Lower mountain coniferous forest (2,650-1,830 meters); restricted to the San Gabriel Mountains
<i>Castilleja grisea</i> San Clemente Island paintbrush	FE, SE, 1B	Coastal scrub, coastal bluff scrub (5-535 meters)
<i>Cercocarpus traskiae</i> Catalina Island mountain-mahogany	FE, SE, 1B	Chaparral, coastal scrub (100-250 meters)
<i>Chloropyron maritimum</i> ssp. <i>maritimum</i> Salt marsh bird's-beak	FE, SE, 1B	Coastal salt marsh, coastal dunes (0-30 meters)
<i>Chorizanthe parryi</i> var. <i>fernandia</i> San Fernando Valley spineflower	FC, SE, 1B	Sandy soils in coastal scrub (3-1,035 meters)
<i>Deinandra minthornii</i> Santa Susana tarplant	1B	Sandstone outcrops and crevices in chaparral and coastal scrub (280-760 meters)
<i>Delphinium variegatum</i> ssp. <i>kinkiense</i> San Clemente Island larkspur	FE, SE, 1B	Valley and foothill grassland (75-500 meters)
<i>Dithyrea maritime</i> Beach spectaclepod	ST, 1B	Coastal dunes, coastal scrub (3-50 meters)

Scientific Name Common Name	Special Status	Preferred Habitat
<i>Dodecahema leptoceras</i> Slender-horned spineflower	FE, SE, 1B	Chaparral, coastal scrub (200□760 meters)
<i>Dudleya cymosa ssp. agourensis</i> Agoura Hills dudleya	FT, 1B	Chaparral and cismontane woodland (200□500 meters)
<i>Dudleya cymosa ssp. marcescens</i> Marcescent dudleya	FT, 1B	Sheer rock faces and rocky cliffs in chaparral (180□520 meters)
<i>Dudleya cymosa ssp. ovatifolia</i> Santa Monica dudleya	FT, 1B	Primarily north-facing slopes with chaparral and coastal scrub (210□500 meters)
<i>Galium catalinense ssp. acrispum</i> San Clemente Island bedstraw	SE, 1B	Steep cliffs and canyons supporting valley and foothill grasslands (20□425 meters)
<i>Helianthemum greenei</i> Island rush-rose	FT, 1B	Chaparral, coastal scrub, closed-cone coniferous forest (15□48 0 meters)
<i>Lithophragma maximum</i> San Clemente Island woodland star	FE, SE, 1B	Moist areas in coastal bluff scrub and coastal scrub (120□400 meters)
<i>Malacothamnus clementinus</i> San Clemente Island bush-mallow	FE, SE, 1B	Valley and foothill grassland (5□275 meters)
<i>Nasturtium gambelii</i> Gambel's water cress	FE, ST, 1B	Marshes and swamps (5□1,305 meters)
<i>Navarretia fossalis</i> Moran's nosegay	FT, 1B	Vernal pools, chenopod scrub, marshes and swamps, playas (30□1,300 meters)
<i>Orcuttia californica</i> California Orcutt grass	FE, SE, 1B	Vernal pools (15□660 meters)
<i>Pentachaeta lyonii</i> Lyon's pantachaeta	FE, SE, 1B	Chaparral and valley and foothill grassland (30□630 meters)
<i>Sibara filifolia</i> Santa Cruz Island rock cress	FE, 1B	Coastal scrub (15□600 meters)

Status Definitions:

USFWS

FE: Species designated as endangered under the federal ESA

FT: Species designated as threatened under the federal ESA

FP: Species designated as protected under the federal ESA

FC: Species is a candidate for listing under the federal ESA

CDFG

SE: Species designated as endangered under the California ESA

ST: Species designated as threatened under the California ESA

SC: Species of Special Concern

CNPS

1B: Plants rare, threatened, or endangered in California and elsewhere

2: Plants rare, threatened, or endangered in California but more common elsewhere

4: Plants of limited distribution

LISTED WILDLIFE SPECIES WITH POTENTIAL TO OCCUR IN THE COUNTY OF LOS ANGELES (CDFG 2010)

Scientific Name Common Name	Special Status ¹	Preferred Habitat
<i>Ammospermophilus nelsoni</i> Nelson's antelope squirrel	ST	Western San Joaquin Valley from 200□1,200 feet above mean sea level on dry, sparsely vegetated loam soils
<i>Anaxyrus californicus</i> Arroyo toad	FE, SC	Semi-arid regions near washes or intermittent streams
<i>Buteo swainsoni</i> Swainson's hawk	ST	Breeds in grasslands with scattered trees; riparian areas, grasslands, and agricultural areas
<i>Catostomus santaanae</i> Santa Ana sucker	FT, SC	Coastal streams
<i>Charadrius alexandrinus nivosus</i> Western snowy plover	FT, SC	Sandy beaches; nests in sandy, gravelly or friable soils
<i>Chelonia mydas</i> Green turtle	FT	Marine environments with adequate supplies of seagrasses and algae
<i>Coccyzus americanus occidentalis</i> Western yellow-billed cuckoo	FC, SE	Nests in riparian forests
<i>Dipodomys merriami parvus</i> San Bernardino kangaroo rat	FE, SC	Sandy loam substrates with alluvial scrub vegetation
<i>Empidonax traillii extimus</i> Southwestern willow flycatcher	FE, SE	Riparian woodlands in southern California
<i>Eucyclogobius newberryi</i> Tidewater goby	FE, SC	Brackish water habitats along the California coast (San Diego County north to the Smith River)
<i>Euphilotes battoides allyni</i> El Segundo blue butterfly	FE	Restricted to remnant coastal dune habitat in southern California
<i>Gasterosteus aculeatus williamsoni</i> Unarmored threespine stickleback	FE, SE (FP)	Small southern California streams with cool, clear water and abundant vegetation
<i>Glaucopsyche lygdamus palosverdesensis</i> Palos Verdes blue butterfly	FE	Palos Verdes Hills in Los Angeles County that support <i>Astragalus tricopodus</i> var. <i>lonchus</i> , its host plant
<i>Gopherus agassizii</i> Desert tortoise	FT, ST	Desert scrub, desert wash, and Joshua tree habitats with friable soils for burrowing and nesting
<i>Gymnogyps californianus</i> California condor	FE, SE	Large areas of grasslands and foothill chaparral in moderate altitude mountain ranges; deep canyons with clefts in rock walls for nesting
<i>Haliaeetus leucocephalus</i> Bald eagle	SE, (FP)	Ocean shore, lake margins, and rivers for nesting and wintering
<i>Laterallus jamaicensis coturniculus</i> California black rail	SE, FP	Freshwater marsh, wet meadows, and shallow margins of saltwater marshes adjacent to larger bays

Scientific Name Common Name	Special Status ¹	Preferred Habitat
<i>Oncorhynchus mykiss irideus</i> Southern steelhead □ southern California DPS	FE, SC	Found from Santa Maria River south to the southern extent of its range in San Diego County
<i>Passerculus sandwichensis beldingi</i> Belding's savannah sparrow	SE	Coastal salt marshes from San Diego County north to Santa Barbara
<i>Perognathus longimembris pacificus</i> Pacific pocket mouse	FE, SC	Narrow coastal plains from the Mexican border north to Los Angeles County; prefers fine alluvial sands
<i>Poliophtila californica californica</i> Coastal California gnatcatcher	FT, SC	Coastal sage scrub
<i>Rana draytonii</i> California red-legged frog	FT, SC	Permanent sources of deep water with dense or emergent riparian vegetation
<i>Rana muscosa</i> Sierra Madre yellow-legged frog	FE, SC	Very near to water in the San Gabriel, San Jacinto, and San Bernardino Mountains
<i>Siphateles bicolor mohavensis</i> Mohave tui chub	FE, SE, FP	Endemic to Mojave River basin; deep pools, ponds, or slough-like areas
<i>Sternula antillarum browni</i> California least tern	FE, SE, FP	Nesting occurs along the coast from the San Francisco Bay south to Northern Baja California
<i>Vireo bellii pusillus</i> Least Bell's vireo	FE, SE	Riparian areas in the vicinity of water or in dry river bottoms below 2,000 feet AMSL
<i>Xerospermophilus mohavensis</i> Mohave ground squirrel	ST	Open desert scrub, alkali scrub, and Joshua tree woodland